

MA 17: HOW TO SOLVE IT
HANDOUT 5: DISCUSSIONS AFTER MOCK 2

Instructor Lingfu Zhang (lingfuz@caltech) **Office** Linde Hall 358

TA Minghao Pan (mpan2@caltech)

Elements in the theory of matrix: trace, determinant, eigenvalues, eigenvectors, characteristic equation/polynomial.

The Cayley–Hamilton Theorem. Any $n \times n$ matrix A satisfies its characteristic equation, which means that if $P_A(\lambda) = \det(\lambda I_n - A)$, then $P_A(A) = O_n$.

The Perron–Frobenius Theorem. Any square matrix with positive entries has a unique eigenvector with positive entries (up to multiplication by a positive scalar), and the corresponding eigenvalue has multiplicity one and is strictly greater than the absolute value of any other eigenvalue.

Some more problems:

Problem 1. (Putnam 1986, B6)

Suppose that A, B, C, D are $n \times n$ matrices (with entries in some field), such that AB^T and CD^T are symmetric, and $AD^T - BC^T = I$. Prove that $A^T D - C^T B = I$.

Hint: find a more “matricial” interpretation of the condition $AD^T - BC^T = I$.

Problem 2. (Putnam 1995, A5)

Let x_1, x_2, \dots, x_n be differentiable (real-valued) functions of a single variable t which satisfy

$$\begin{aligned} \frac{dx_1}{dt} &= a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ \frac{dx_2}{dt} &= a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ &\vdots \\ \frac{dx_n}{dt} &= a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n \end{aligned}$$

for some constants $a_{ij} > 0$. Suppose that for all i , $x_i(t) \rightarrow 0$ as $t \rightarrow \infty$. Are the functions x_1, x_2, \dots, x_n necessarily linearly dependent?

Problem 3. (Putnam 1999, B5)

For an integer $n \geq 3$, let $\theta = 2\pi/n$. Evaluate the determinant of the $n \times n$ matrix $I + A$, where I is the $n \times n$ identity matrix and $A = (a_{jk})$ has entries $a_{jk} = \cos(j\theta + k\theta)$ for all j, k .

Hint: find the eigenvectors, and use complex numbers.

Another problem on complex numbers:

Problem 4. (Putnam 1991, B2)

Suppose f and g are non-constant, differentiable, real-valued functions defined on $(-\infty, \infty)$. Furthermore, suppose that for each pair of real numbers x and y ,

$$f(x+y) = f(x)f(y) - g(x)g(y),$$

$$g(x+y) = f(x)g(y) + g(x)f(y).$$

If $f'(0) = 0$, prove that $(f(x))^2 + (g(x))^2 = 1$ for all x .

Reminder PSet 5 to be released tomorrow, and due by the end of Nov 24 (on Canvas). The fourth problem session (to discuss PSet 4) will be on Nov 13, 7-8 PM, at the same location.